## INDEX TO VOLUME 4.

Names of new plants are printed in full face type.

Abies balsamea, 24, 84, 189.

Acer dasycarpum, 130; Negundo, 130;

saccharinum, 130. Acerates viridiflora, 17.

Acolea coralloides, 207.

Acrosiphonia, 112, 115.

Actaea, 88; alba, 75. Adder's Tongue, 88.

Additions to the Flora of Massachusetts, Some, 166.

Adiantum pedatum, 247.

Adpressum Group, Two New Hyperi-

cums of the, 135. Aegagropila, 113, 115.

Agarics, Pink-spored, 6.

Agaricus campestris, 77, 79, 199.

Agrimonia gryposepala, 91; parviflora, 85; striata, 91.

Agrostis, 49; alba, 33; arachnoidea, 143; brevifolia, 46, 47; coarctata, 26; intermedia, 39, 90; juncea, 48, 49; perennans, 90; rubicunda, 143; tri-

chopodes, 143. Aira artica, 143; brevifolia, 143, 144;

media, 144.

Ajuga reptans, 14. Alder, 74, 168.

Algae, 111, 118, 124.

Algologist's Vacation in Eastern Maine,

Alnus viridis, 189.

Amanita, 5, 6, 78; caesarea, 5; muscaria, 78.

Amanitopsis, 5, 6; vaginata, 78.

Amblystegium, 238, 239; adnatum, 238; chrysophyllum, 238; compactum, 239; hispidulum, 239; irriguum, 182, 239; Lescurii, 182; serpens, 239; subtile, 181.

America, An Osmorrhiza New to Eastern, 153; Early Records of Leontodon in, 39; Taraxacum palustre in, 155; Vari-

ations of Glaux in, 213.

American Cranberries, Variations and Distribution of, 231; Horticulture, Cyclopedia of, 108; Occurrence of the European Centaurea diffusa, 249; Variety of Vaccinium Vitis-Idaea, 231.

Ammidown, L. E., Saleratus as a Protection against Rhus-poisoning, 105.

Ammophila, 33. Amphicarpa Pitcheri, 91.

Anacamptodon splachnoides, 239.

Anadema, 115.

Anagallis arvensis, 16, 185, 186, and caerulea in Vermont, 185; caerulea 185, 186.

Andover, Massachusetts, Cuphea procumbens at, 247; Hieracium praealtum

Andrews, A. L., List of Bryophytes from the Mt. Greylock Region, 29; Bryophytes of the Mt. Greylock Re-gion,—II, 238; Note upon Recent Treatment of Habenaria hyperborea

and its Allies, 79.

Andrews, L., Gleditschia triacanthos established in Connecticut, 103;

Notice of work, 202. Andromeda ligustrina, 131.

Anemone, 88; quinquefolia, 88; virginiana, 74.

Anemonella thalictroides, 63, 76.

Aneura latifrons, 242.

Annual Meeting of the Josselyn Society, 188; Winter Meeting of the Vermont Botanical Club, Seventh, 64.

Anomalous Fern, 10; Skullcap, 137.

Anomodon apiculatus, 30; attenuatus, 30; rostratus, 239.

Antennaria canadensis, 85, 92; neodioica, 92; Parlinii, 152, Staminate Plant of, 152; plantaginea, var. petiolata, 92. Anthelia julacea, 213; Juratzkana, 212,

213.

Antithamnion boreale, 177; Pylaisaei, 177.

Apargia autumnalis, 39.

Aphyllon uniflorum, 76. Aplectrum hiemale, 18, 20, 22.

Apocynaceae, 17. Apocynum, 20; androsaemifolium, 17, 81; cannabinum, 17, 81; medium,

Aquilegia canadensis, var. flaviflora, 169, in Maine, 169, var. Phippenii, 169.

Arabis confinis, 85; perfoliata, 93.

Aralia racemosa, 75. Arbutus, 127, 128; Trailing, 140.

Arceuthobium, 249; pusillum, 28, 249; Two New Stations for, 249.

Arctostaphylos, 34. Arenaria peploides, 189. Arethusa bulbosa, 18, 76. Armillaria, 77; mellea, 77.

Arum, 168.

Asclepiadaceae, 17.

Asclepias Cornuti, 17; incarnata, 17, var. pulchra, 17; obtusifolia, 17, 19; phytolaccoides, 17; purpurascens, 17; quadrifolia, 17, 19, 74; syriaca, 17; tuberosa, 17, 19; verticillata, 17, 19.

Ash, Prickly, 130.

Aspideae, 9; New England, 7.

Aspidium, 7-9, 51, 52, 53, 160, 164, 166; aculeatum, var. Braunii, 247; Braunii, 10; Goldianum, 247; Lonchitis, 8; Plumieri, 160; spinulosum, var. dilatatum, 247, var. intermedium, 247; trifoliatum, 8, 160.

Asplenium, 165; acrostichoides, ebenoides, 54; ebeneum, 54, var. Hortonae, 54; Filix-foemina, 7; montanum, 54, 157; Ruta-muraria, 55; Trichomanes, 54; viride, 54.

Asprella Hystrix, 89.

Aster, 89, 186; cordifolius, 88; ianthinus, 203; macrophyllus, var. ianthinus, 203; Novae-Angliae, 89; Novi-Belgii, 92, 186; salicifolius, 89, 94; tardiflorus, 100; undulatus, 186; undulatus X Novi-Belgii, 186, 187.

Athyrium, 8, 160; Filix-foemina, 160. Atrichum angustatum, 30; undulatum,

Atriplex patulum, 92.

Aulacomnium heterostichum, 30, 169, in Maine, 169.

Avena striata, 99.

Avicularia, 65, 66; New England Polygonums of the Section, 65.

Azalea, 141; viscosa, 106.

Bachbunge, 191.

Bacon, A. E., Anagallis arvensis and caerulea in Vermont, 185; Cases of Poisoning by Cypripedium spectabile in Vermont, 94; Dwarf Mistletoe at Bradford, Vermont, 28.

Bailey, L. H., Notice of work, 108. Bailey, W. W., Some Notable Plant Stations in Rhode Island, 198.

Barbarea vulgaris, 92.

Barbula unguiculata, 30.

Bartramia pomiformis, 30. Bazzania trilobata, 30.

Beaked Hazel, 132. Beal, W. J., Seed-throwing of Viola, 230.

Beccabunga, 191.

Beech, 93, 169; Blue, 132.

Betula lenta, 83; papyrifera, 90; pendula, var. dalecarlica, 83; populifolia, 132.

Bidens vulgata, 100.

Bigelow, E. G., Notice of work, 40. Biological Relation of Polygonum Hartwrightii to P. amphibium, 104.

Birch, 74, 141; A Cut-leaved Cherry, 83; Cherry, 83; Gray, 132; White, 83.

Bird Club, The Vermont, 138.

Bird's Knotgrass, 65.

Bissell, C. H., Biological Relation of Polygonum Hartwrightii and P. amphibium, 104; Newly Introduced Species of Crepis and Leontodon, 249; Notice of work, 202; Rediscovery of Phaseolus perennis in New England, 13; Some Noteworthy Plants of Connecticut, 98. Bitter Boletus, A Form of, 187,

Black Chokeberry, 55, 56; Huckleberry, 14; Oak, 132, Scrub, 133; Spruce, 35.

Blackberry, 40. Bladder-nut, 130.

Blanchard, W. H., Elusive Character of Pogonia pendula, 2; Our Chokeberries, 55; Some Vermont and New Hampshire Plants in the Middle Connecticut Valley, - I, 129.

Blasia pusilla, 30.

Blepharostoma trchophyllum, 30.

Blomberg, C., Some Plants recently found in and around North Easton, Massachusetts, 13; Utricularia minor in Vinnicunnet, Massachusetts, 102.

Bloodroot, 128.

Blooming of Hepaticas, 127.

Blue Beech, 132; Hills Reservation, A Few Plants of the, 74.

Blue-berried Huckleberry, Notes on the,

Blueberry, 14; Half-high, 131.

Blue-black Huckleberry, 14.

Boleti, 172.

Boletus, 78; A Form of the Bitter, 187;

felleus, 187, 188, var. obesus, 187. Botanical Club, The Vermont, Seventh Annual Winter Meeting of, 64; Society of Maine, Josselyn, 110, 188, Mid-winter Meeting, 62.

Botany, Outlines of, 85.

Botrychium, 159; lanceolatum, 36; matricariaefolium, 36; ternatum, 55.

Botrypus, 159. Box Elder, 130.

Brachythecium glaciale, 182; reflexum, 30, 182; salebrosum, 30; Starkei, 30. Bradford, Vermont, Dwarf Mistletoe at, 28.

Brainerd, E., Two More Rare Plants from Lake St. John, Quebec, 128.

Brasenia peltata, 76.

Brassica campestris, 92; juncea, 92; Sinapistrum, 92.

Broklempe, 191.

Broklympe, 191.

Bromus brachyphyllus, 146; laevipes, 147; secalinus, 90.

Brooklime, 81, 191, 192.

Brooks, H., Notice of work, 40.

Brunswick, Maine, Occurrence of Eleocharis diandra at, 1.

Bryhnia graminicolor, 183.

Bryophytes, 29; from the Mt. Greylock Region, List of, 29; of the Mt. Greylock Region, — II, 238.

Bryum affine, 181; argenteum, 30; bimum, 30; caespiticium, 30; pseudotriquetrum, 181; roseum, 30.

Buffalo Bur, 151. Bulrush, Giant, 102. Bur, Buffalo, 151. Buttercup, 32.

Button bush, 131, 243, 244; Remarkable Persistence of, 243.

Buttonwood, 132.

Buxbaumia aphylla, 239; indusiata, 239.

Calamintha Acinos, 246; Clinopodium, 246.

Calla palustris, 74, 106. Callitriche Austini, 99.

Calluna vulgaris, 200; in Southern New Hampshire, 200.

Calopogon pulchellus, 18. Calothrix pulvinata, 178. Calypso borealis, 18, 22.

Campanula aparinoides, 15; glomerata, 15, 20; rapunculoides, 17; rotundifolia, 15, var. arctica, 19.

Campanulaceae, 15.

Camptosorus, 63; rhizophyllus, 55, 63. Cardamine hirsuta, var. sylvatica, 92; pennsylvanica, 92.

Carduus discolor, 203.

Carex, 167; acutiformis, 218; adusta, 218; aenea, 218, 227; aestivalis, 218, 227; alata, 218; albicans, 218, 223, 224; albolutescens, 218, 227; alopecoidea, 218; alpina, 228; aperta, 226; aquatilis, 218, var. elatior, 218; arcta, 218; arctata, 218, var. Faxoni, 218, X castanea, 218; atrata, var. ovata, 218; atlantica, 229; aurea, 168, 218, in Connecticut, 168; Backii, 218, Bebbii, 218, 228; Bicknellii, 219, 228; bromoides, 219; brunnescens, 219; bullata, 219, var. Olneyi, 219; canescens, 219, 228, var. disjuncta, 219, var. subloliacea, 219; capillaris, 219, var. elongata, 219, 228; capitata, 150, 219, 228; Careyana, 227; castanea, 219, 228; cephaloidea, 219; cephalophora, 219; chordorrhiza, 219, 228; concinna, 228; conoidea, 219; Crawei, 219, 227, 228; Crawfordii, 219, var. vigens, 219; crinita, 219, var. gynandra, 219, var. minor, 219, var. simulans, 219, X torta, 219; cristata, 219; Davisii, 219; debilis, var. Rudgei, 219, var. strictior. 219, X virescens, 219; deflexa, 219, var. Deanei, 219; Deweyana, 219; digitalis, 219; divulsa, 225; eburnea, 219, 228; echinata, 222, 225, var. cephalantha, 222, var. ormantha, 222; elachycarpa, 227; Emmonsii, 223, var. elliptica, 223, 224; exilis, 219, 229; festucacea, 219, var. brevior, 219; filiformis, 219, 224, var. latifolia, 219; flava, 219, 224, var. Oederi, 219, 224; foenea, 219, var. perplexa. 219; folliculata, 219; formosa, 220; fulva, 227; fusca, 220; glaucodea, 60, 220; Goodenowii, 220, 224, 225; gracillima, 220; Grahami, 220; granularis, 220, var. Haleana, 220; Grayi, 220, 229; grisea, 220, var. angustifolia, 220; gynocrates, 220, 227, 229; Haydeni, 226; hirta, 220; Hitchcockiana, 220; Houghtonii, 220; hystricina, 220, var. Dudleyi, 220; interior, 220, var. capillacea, 220; intumescens, 37, 220, var. Fernaldii, 220; katahdinensis, 128, 220; laevigata, 227; laxiculmis, 220; laxiflora, 220, var. latifolia, 220, var. patulifolia, 220, var. striatula, 220, var. varians, 220; lenticularis, 220; leporina, 220, 229; leptalea, 220; limosa, 220, 224; littoralis, 99, 220; livida, 220; longirostris, 220; lupulina, 220, var. pedunculata, 220, var. polystachya, 220, X bullata, 220; lurida, 220, var. exudans, 220, var, flaccida, 220, var. gracilis, 220, 229, var. parvula, 220, X lupulina, 221; magellanica, 150, 221, 229; maritima, 221, var. erectiuscula, 221; Michauxiana, 221; mirabilis, 37, 221, var. perlonga, 90, 221, var. tincta, 221; Muhlenbergii, 221, var. enervis, 37,

Q. 221; muricata, 221; nigro-marginata, 27, 221; norvegica, 221; Novae-Angliae, 221, 229; Oederi, 224; oligocarpa, 221; oligosperma, 221, 229; oronensis, 221; pallescens, 221; panicea, 221; pauciflora, 221; peduncu- Cariceae, 227. lata, 84, 221, 229; pennsylvanica, 221, Carices at Suc var. lucorum, 221; pilulifera, 221, var. longibracteata, 221; plantaginea, 221; platyphylla, 221; polymorpha, 221; 229; Porteri, 221; praecox, 221; prasina, 221; pratensis, 221, 224; praticola, 224; Pseudo-Cyperus, 221, var. americana, 221; ptychocarpa, 221; pubescens, 221; rariflora, 221; retrorsa, 221, var. Hartii, 221; retrorsa X rostrata, 221, X utriculata, 221; rigida, 224, 225, var. Bigelovii, 221, 225, var. Goodenovii, 224; riparia, 221; rosea, Cedar, Red, 133. 221, var. minor, 222. var. radiata, 222, 227, var. retroflexa, 222; rostrata, Centaurea diffusa, 249, American Occur-222, var. ambigens, 222, var. utricucuspidata, 222; saxatilis, 222, var. milia-225; scirpoidea, 222, 229, 250; scoparia, 222, var. condensa, 222, var. monili- Ceramium rubrum, 177. formis, 222; seorsa, 222; setacea, Cerastium arvense, var. fuegianum, 150. 222; Shortiana, 227; siccata, 222; Ceratodon purpureus, 30. silicea, 33, 222; sparganoides, 222; Certain Eaters of Mushrooms, 77. squarrosa, 222, 225; stellulata, 222, Cesia coralloides, 207. 225, var. angustata, 222, var. cephalan- Chamaecyparis sphaeroidea, 106. ormantha, 222; sterilis, 222, 229. var. excelsior, 222; stipata, 222; Character of Pogonia pendula, The straminea, 222; striata, 227, var brevis, 222, 230; stricta, 222, 225, Cheney, C. I., Rare Plants in Centre-226, var. angustata. 222, 226. var. cur-ville, Massachusetts. 245. tissima, 222, 226, var. decora, 222, Cherry, 81; Birch, 83, Cut-leaved, 83; 226,  $\times$  salina, 222,  $\times$  torta, 222; sty-Sand, 130. loflexa, 222; subulata, 223, 230; Chervil, 81. tenella, 223; tenera, 223, var. invisa, Chestnut, 133. 223, var. Richii, 223; tenuiflora, 223, Chilian Empetrum in New England, 147. 226, 230; teretiuscula, 223, 230, var. Chiloscyphus polyanthos, 30. ramosa, 167, 223; tetanica, 168, 223, Chimaphila maculata, 75. var. Meadii, 223, 227, var. Woodii, Chlorophyceae, 178. 167, 223; torta, 223; tribuloides, 223, Chokeberries, Our, 55. var. reducta, 223. var. turbata, 223; Chokeberry, Black, 55, 56, Red, 55, 131. triceps, var. hirsuta, 223; trichocarpa, Chorda Filum, 175; tomentosum, 175. 223, 227, 230; trisperma, 223, 226; Christmas Fern, 10, 53. trisperma × tenuiflora, 223, 226; Chrysosplenium, 250; americanum, 76. 225; umbellata, 223, var. brevirostris, 223, var. tonsa, 223; vaginata, 223; Circaea intermedia, 137; Lutetiana, 89. jejuna, 223, var. monile, 223; vestita, tive of New Hampshire, Is, 217.

223, 227, var. Kennedyi, 223; virescens, 223, var. costata, 223, X arctata, 223; vulgaris, 224; vulpinoidea, 223; Willdenovii, 223; xanthocarpa, 223, var. annectans, 223.

Carices at Sudbury, Massachusetts, Two Noteworthy, 167.

Carpinus caroliniana, 132.

Carval, 81.

Carvell, 81.

Carya alba, 132; amara, 132.

Cases of Poisoning by Cypripedium spectabile in Vermont, 94.

Cassia Chamaechrista, 37.

Castanea sativa, var. americana, 133.

Cat-tail Flag, 93.

Ceanothus americanus, 76, 130.

Celtis occidentalis, 132.

rence of the European, 249; nigra, 189. lata, 222; rotundata, 222; salina, var. Centreville, Massachusetts, Rare Plants in, 245.

ris, 222, var. rhomalea, 222; scabrata, Cephalanthus, 182, occidentalis, 131, 243. 222, X crinita, 222; Schweinitzii, 222, Cephalozia curvifolia, 242; divaricata, 242; multiflora, 242.

tha, 222, var. excelsior, 222, var. Chamberlain, E. B., Aulacomnium heterostichum in Maine, 169.

Tuckermani, 223; typhinoides, 223, Churchill, J. R., Some Plants from Prince

Edward Island, 31.

varia, 223, 224, 227, var. colorata, 223; Cirsium altissimum, var. discolor, 203; vesicaria, 223, var. distenta, 223, var. horridulum, 93; palustre, 217, a NaCistaceae, 28.

Cladophora, 111, 112, 113, 115, 124, 125; albida, 114, 119, 120, 121, 125, 127, var. refracta, 119; arcta, 114, 115, 116, 117, 118, 127, var. centralis, 117; diffusa, 126; expansa, 114, 122, 123, 124, 127, 178, var. glomerata, 123; flavescens, 124; flexuosa, 114, 119, 121, 122, 125, 127, forma densa, 121; fracta, 114, 124, 127, forma flavescens, 124, forma marina, 124; gracilis, 114, 120, 121, 122, 123, 126, 127, forma elongata, 122, var. expansa, 122, var. tenuis, 122, 123, var. vadorum, 122; glaucescens, 114, 120, 121, 127; hamosa, var. refracta, 125; hirta, 114, 121, 122, 123, 126, 127; Hutchinsiae, 114, 126, 127, var. distans, 126; Hystrix, 114, 116, 117, 127; lanosa, 114, 118, 119, var. uncialis, 118; laetevirens, 114, 121, 122, 125, 127, var. glomerata, 126; Magdalenae, 114, 124; refracta, 114, 119, 120, 124, 125, 127; Rudolphiana, 114, 120, 127; rupestris, 114, 125, 127; Sonderi, 116, 117; spinescens, 114, 116, 117, 118, 119, 127; uncialis, 118, utriculosa, var. laetevirens, 126.

Cladophoras of New England, The Marine, III.

Hills Reservation, 74.

brook, Massachusetts, 42.

Clark, H. S., Erodium moschatum in Coreopsis trichosperma, 76. Connecticut, 248.

Clathrus, 134; cancellatus, 135; columnatus, 134, 135, in Lawrence, Massachusetts, 134

Clethra, 108

Cliff-Brake, Purple, 54.

Climacium americanum, 239, var. Kinbergii, 239.

Climbing Fern, 51; in New Hampshire,

Clintonia borealis, 74, 76, 106.

Clitocybe, 7.

Club, Field Meeting of the Vermont Botanical, 189; The Vermont Bird, 138, The Vermont Botanical, 138.

Club Moss, 55.

Cnicus horridulus, 93; lanceolatus, 92.

Coast at Cutler, Maine, 23.

Collins, F. S., Algologist's Vacation in Eastern Maine, 174; Marine Cladophoras of New England, 111.

Collins, J. F., Distinctive Features of Cryptotaenia, 88. for the Typical Lycopodium complanatum, 154.

Collinsonia canadensis, 89. Colpodium junceum, 48.

Coltsfoot, Sweet, 128.

Comandra livida, 18, 98, and Lycopodium sitchense on Mt. Washington, Discovery of, 97; umbellata, 18.

Compositae, 33.

Concerning the Plants mentioned in

Young's Chronicles, 81.

Conferva, 112; albida, 119; arcta, 116; diffusa, 216; distans, 126; expansa, 123; flexuosa, 121; glaucescens, 120; gracilis, 121; Hutchinsiae, 126; laetevirens, 126; lanosa, 118; refracta, 119, 120, 125; Rudolphiana, 120; rupestris, 125; uncialis, 118; vadorum, 122.

Connecticut, Carex aurea in, 168; Erodium moschatum in, 248; Gleditschia triacanthos established in, 103; Notes on the Flora of, 36; Noteworthy Plants of, 84, of Southeastern, —III, 26; Some Noteworthy Plants of, 98; Valley, Some Vermont and New Hampshire Plants in the Middle, - I, 129.

Conocephalus conicus, 30.

Conopholis, 169; americana, 75, 76, 169, in Maine, New Stations for Peltandra virginica and, 168.

Clark, A., A Few Plants of the Blue Coprinus atramentarius, 199; comatus, 199.

Clark, A. G., Utricularia minor in Hol- Corallorhiza innata, 18, 74; multiflora, 18, 75; odontorhiza, 18; striata, 22.

Cornus, 89; canadensis, 74, 75; circinata,

131; paniculata, 89, 131. Corylus americana, 132; rostrata, 132. Cotton-grass new to North America, 82.

Cotula coronopifolia, 32, 33.

Cowberry, 232.

Cow Lily, 93. Cranberries, Variations and Distribution

of American, 231.

Cranberry, High, 131; Large, 234; Mountain, 231, 232; Rock, 231; Small, 235. Crataegus, 64; asperifolia, 166, 167; lobu-

lata, 167; Pringlei, 167; scabrida, 166. Crepis, 249; and Leontodon, Newly Introduced Species of, 249; rigida, 250; taraxacifolia, 250; virens, 197, in Eastern Massachusetts, 197.

Cress, Garden, 32; Water, 81.

Cristate Form of Nephrodium marginale. 244.

Crowberry, 34, 147.

Iris Hookeri, 179; Extension of Range Cuphea, 247; procumbens, 248, at Andover, Massachusetts, 247: viscosissima.

Cuscuta Gronovii, 92. Cut-leaved Cherry Birch, 83. Cutler, Maine Coast at, 23. Cyanophyceae, 178.

Cyclopedia of American Horticulture,

108.

Cylindrothecium cladorhizans, 240; seductrix, 240.

Cyperaceae, 82, 225.

Cypripedium, 22, 95, 97; acaule, 18, 106; arietinum, 18, 20, 62, on Mt. Toby, Massachusetts, 62; parviflorum, 18, 21; pubescens, 18, 96; spectabile, 18, 95, 96, 97, in Vermont, Some Cases of Poisoning by, 94.

Cystopteris bulbifera, 247; fragilis, 150,

Dactylis glomerata, 90. Dame, L. L., Notice of work, 40. Dandelion, 155; Fall, 39. Davenport, G. E., Miscellaneous Notes on New England Ferns, - III, 7, IV, 49, V, 157.

Davis, C. A., Occurrence of Eleocharis

diandra at Brunswick, Maine, 1. Deane, W., Calluna vulgaris in Southern New Hampshire, 200; Cyclopedia of American Horticulture, 108; Remarkable Persistence of the Button-bush,

243. Delesseria alata, 176.

Deschampsia arctica, 143, 144; brachyphylla, 143, 144; brevifolia, 143, 144; caespitosa, 144, var. alpina, 144, var. arctica, 144, var. brevifolia, 144, var. setifolia, 144; curtifolia, 143, 144; setifolia, 144.

Desmodium acuminatum, 89; canadense, 37; canescens, 37; cuspidatum, 37, 90; humifusum, 27; paniculatum, 75;

rotundifolium, 75. Devil's Paint-brush, 62. Diapensia lapponica, 15.

Diapensiaceae, 15.

Dichelyma capillacea, 182.

Dicranella heteromalla, 30; varia, 30. Dicranum flagellare, 240; fulvum, 240; fuscescens, 30, 242; longifolium, 30; montanum, 182; scoparium, 30, 240,

var. pallidum, 240. Dictyophora, 134. Diervilla trifida, 75. Diospyros virginiana, 16, 19.

Diphyscium foliosum, 30.

Dirca palustris, 132.

Discovery of Comandra livida

Lycopodium sitchense on Mt. Washington, 97.

Distinctive Features of Iris Hookeri, 179. Distribution of American Cranberries, Variations, and, 231.

Ditrichum tortile, 240; vaginans, 181,

240. Dogwood, Poison, 106; Round-leaved,

131. Draba caroliniana, 85; incana, 150.

Driggs, A. W., Notes on the Flora of Connecticut, 36.

Drosera intermedia, 42; rotundifolia, 42. Drummondia clavellata, 30.

Druopteris, 163.

Dryas octopetala, 98.

Dryopteris, 10, 163, 164, 166; Filix-mas,

164; simulata, 64. Dwarf Mistletoe, 35, at Bradford, Vermont, 28; Sumach, 130; Tansy, 32.

Early Records of Leontodon in America,

Eastern America, An Osmorrhiza New to, 153; Maine, Algologist's Vacation in, 174; Massachusetts Crepis virens in, 197, Hemicarpha in, 82, Plants new to, 60.

Eaters of Mushrooms, Certain, 77.

Ebenaceae, 16.

Ebony Spleenwort, 54.

Ectocarpus tomentosus, 118.

Eggleston, W. W., Discovery of Comandra livida and Lycopodium sitchense on Mt. Washington, 97; Stature attained by Euphrasia Oakesii, 108.

Elder, 108; Box, 130. Eleocharis diandra, 2, at Brunswick, Maine, Occurrence of, 1; palustris, var. vigens, 27.

Ellisia Nyctelea, 170, in Massachusetts,

Juncus Torreyi and, 170.

Elusive Character of Pogonia pendula, 2. Elwell, L. H., Cypripedium arietinum on Mt. Toby, Massachusetts, 62.

Elymus aristatus, 147; hirsutiglumis, 27; robustus, 27; triticoides, 147.

Empetrum, 34, 196; in Franklin County, Maine, Occurrence of, 196; in New England, The Chilian, 147; nigrum, 24, 147, 148, 150, 189, 196; var. andinum, 148, 149, 150, 196; purpureum, 148, 149; rubrum, 148, 149.

Enteromorpha, 123, 175; intestinalis, 175.

Entodon repens, 183. Epipactis Helleborine, 22. Epiphegus virginiana, 93.

and Equisetum hiemale, 55.

Eragrostis campestris, 144; major, 39; nitida, 144; Purshii, 39; reptans, 39. Ericaceae, 200.

Eriocaulon septangulare, 103.

Eriophorum, 82; polystachyon, 82, var. Vaillantii, 82; virginicum, 82.

Erodium, 248; moschatum, 248, in Connecticut, 248.

Errata, 250.

Erythronium, 127, 128; americanum, 88; Yellow, 128.

Eucladophora, 113, 114, 115, 119.

Eunephrodium, 162.

Eupatorium aromaticum, 76; purpureum,

Euphorbia corollata, 85.

Euphrasia, 108; americana, 26, 189; Oakesii, 108, Stature attained by, 108; Randii, 26.

Eurhynchium strigosum, 240, var. robustum, 183.

Evans, A. W., Notes on New England Hepaticae, 207.

Evergreen Ferns in New England, 49.

Everlastings, 34. Extension of Range for the Typical Lycopodium complanatum, 154.

Fagopyrum esculentum 203, 206; tataricum, 203, 206.

Fall Dandelion, 39. False Maiden-hair, 54.

Features of Iris Hookeri, Distinctive, 179. Fern, An Anomalous, 10; Christmas, 10, 53; Climbing, 51, in New Hampshire, 83.

Fernald, M. L., Anomalous Skullcap, 137; Aster undulatus X Novi-Belgii, 186; Chilian Empetrum in New England, 147; Cotton-grass new to North America, 82; Early Records of Leontodon in America, 39; Osmorrhiza New to Eastern America, 153; Preliminary Lists of New England Plants,—X, Carex, 218; Seneca Snakeroot in Maine, 133; Taraxacum palustre in America, 155; Two Northeastern Veronicas, 191; Variations and Distribution of American Cranberries, 231; Varieties of Glaux in America, 213.

Ferns, Miscellaneous Notes on New England,—III, 7, IV, 49, V, 157; of Franklin County, Maine, Notes on some, 247; of New England, Evergreen, 49.

Festuca ovina, 33; tenella, 39.

Few Plants of the Blue Hills Reservation, A, 74.

Field Book, New Popular, 201; of American Wild Flowers, 201; Meeting of the Vermont Botanical Club, 189.

Fimbriaria tenella, 30.

Fissidens adiantoides, 30; decipiens, 30; incurvus, 181, 182; minutulus, 182; taxifolius, 240.

Flag, Cat-tail, 93.

Flax, 81.

Fleshy Fungi, 171.

Fletcher, E. F., Lonicera sempervirens established in Westford, Massachusetts, 40.

Flora, Destruction of our Native, 200; New Local, 202; Oak Island and its, 87; of Connecticut, Notes on the, 36; of Massachusetts, Some Additions to the, 166; Protection of Our Native, Floyd, F. G., Cristate Form of Nephro-

dium marginale, 244; Ranunculus abortivus, var. eucyclus in Hyde Park,

Massachusetts, 152.

Fontinalis Cardoti, 182; Lescurii, 182. Form of Nephrodium marginale, Cristate, 244; of the Bitter Boletus, 187.

Fougère mâle, 158, 162.

Fox Grape, 130.

Franklin County, Maine, Notes on some Ferns of, 247; Occurrence of Empetrum in, 196.

Fraxinus americana, 17; pennsylvanica, 17. var. lanceolata, 17, 19; pubescens, 17; sambucifolia, 17; viridis, 17, 19.

Frost Herbaria at the University of Vermont, Pringle and, 171.

Frullania aeolotis, 242; Asagrayana, 30; eboracensis, 30; riparia, 242.

Fuci, 125.

Funaria hygrometrica, 30.

Fungi, 77, 171.
Further Notes on Solanum rostratum and Hieracium praealtum in Maine, 151; Observations on Nephrodium, Some; Stations for Veronica Chaemaedrys, 152.

Galium Aparine, 195; erectum, 91; Mollugo, 91; trifidum, 26.

Garden Cress, 32.

Gaylussacia resinosa, 14, 131, var. glaucocarpa, 14, 131.

Gemmae of Lycopodium lucidulum, 58. Gentiana Andrewsii, 93.

Geranium maculatum, 88; Robertianum,

Gerardia quercifolia, 88. Geum rivale, 76. Giant Bulrush, 102.

Glaux, 213, 214, 215; in America, Variations of, 213; maritima, 16, 33, 214, 215, var. obtusifolia, 215.

Gleditschia triacanthos, 103; established in Connecticut, 103.

Glyceria elongata, 39, 146. Gnaphalium sylvaticum, 34. Golden-rod, Seaside, 89.

Gonatherus, 153.

Goodyera Menziesii, 18; pubescens, 18, 75; repens, 21, var. ophioides, 18, 21; tesselata, 18.

Grape, Fox, 130; Summer, 130.

Grasses, Notes on North American, 142.

Gratiola aurea, 76.
Graves, C. B., Noteworthy Plants of Southeastern Connecticut,—III, 26; Valerianella in New England, 195.

Gray Birch, 132.

Green Spleenwort, 54.

Greylock, Pogonatum capillare on Mt., 83; Region, A List of Bryophytes from the Mt., 29, Bryophytes of the Mt.,- II, 238.

Grimmia apocarpa, 240; maritima, 24. Grout, A. J., Notes on Vermont Mosses, 180.

Gutterson, M. E., Cuphea procumbens at Andover, Massachusetts, 247.

Gymnomitrium, 208; adustum, 208; concinnatum, 207; coralloides, 207, 208.

Habenaria, 74, 79; blepharoglottis, 18, 74; bracteata, 18; ciliaris, 18, 21; dilatata, 18, 80; fimbriata, 18, 106; Hookeriana, 18, var. oblongifolia, 18, 21; hyperborea, 18, 80, and its Allies, Note upon the Recent Treatment of, 79; lacera, 18, 74; obtusata, 18, 22; orbiculata, 18, 22; psychodes, 18, 74; tridentata, 18, 74; virescens, 18.

Hackberry, 132

Half-high Blueberry, 131. Hamamelis virginiana, 96.

Handbook of Trees of New England, 40. Harger, E. B., Noteworthy Plants of Connecticut, 84.

Harper, R. M., Notes on Lycopodium clavatum and its Variety monostachyon, 100.

Harvey, L. H., Further Notes on Solanum rostratum and Hieracium praealtum in Maine, 151.

Hawkweed, Mouse-ear, 151; Orange, 62,

151. Hazel, Beaked, 132.

Hazemut, 132. Hazen, T. E., Seventh Annual Winter Meeting of the Vermont Botanical Club, 64.

Hedwigia ciliata, 30.

Hedysarum cuspidatum, 90.

Helenium nudiflorum, 198. Heleochloa juncea, 48.

Helianthemum majus, 91.

Helianthus divaricatus, 88; strumosus, 88.

Heliopsis laevis, 100.

Hemicarpha, 82, in Eastern Massachusetts, 82; subsquarrosa, 82.

Hemlock, 93, 169.

Hemp, 81.

Hepatica, 127, 128.

Hepaticae, 30, 242; Notes on New England, 207. Hepaticas, Blooming of, 127.

Hepatics, 29. Hervey, E. W., Rare Plants about New Bedford, 106.

Hieracium, 25, 60, 197; aurantiacum, 62, 151; floribundum, 25; Pilosella, 151; praealtum, 61, 62, 151, 197, at Andover, Massachusetts, 197, in Maine, Further Notes on Solanum rostratum and, 151; vulgatum, 60.

High Cranberry, 131.

Hoary Pea, 249.

Hoffmann, R., Virulence of the Wild Parsnip, 188.

Holbrook, Massachusetts, Utricularia minor in, 42.

Holly, 140, 142.

Holt, G. W., Is Cirsium palustre a Native of New Hampshire, 217.

Homalia Jamesii, 183. Honeysuckle, Trumpet, 40.

Hop-hornbeam, 93.

Horsenettle, 13.

Horticulture, Cyclopedia of American, 108.

Hottonia inflata, 16, 20, 76, 85.

Houstonia, 75; caerulea, 75; purpurea, var. longifolia, 75

Huckleberry, 131; Black, 14; Blue-black, 14; Notes on the Blueberried, 14.

Hudsonia ericoides, 61; tomentosa, 34,

Hyde Park, Massachusetts, Ranunculus abortivus, var. eucyclus in, 152. Hydrocotyle umbellata, 103.

Hylocomium brevirostre, 181; pyrena-

icum, 182; splendens, 30; squarrosum, 181; triquetrum, 30; umbratum,

Hypericum, 135; adpressum, 135, 136, 137, var. spongiosum, 136, 137;

Bissellii, 136, 137; ellipticum, 103. Hypericums of the Adpressum Group,

Îwo New, 135.

Hypnum, 238; chrysophyllum, 238; cordifolium, 182; crista-castrensis, 30, 181; cupressiforme, 240; curvifolium, 181, 241; dilatatum, 241; eugyrium, 241; fertile, 181; fluitans, var. atlanticum, 181; Haldanianum, 241; hispidulum, 239; imponens, 181; 241; molluscum, 182; ochraceum, 182; pallescens, 181, 241; Patientiae, 181; pratense, 30, 181; reptile, 181, 241; Schreberi, 30; uncinatum, 30, forma plumosa, 182.

Ilea fulvescens, 175.

Ilex verticillata, var. tenuifolia, 92.

Intervale, New Hampshire, Some plants

Introduced Species of Crepis and Leontodon, Newly, 249.

Iris, 89; Hookeri, 24, 179, 180, Distinctive Features of, 179; prismatica, 89;

versicolor, 24, 179, 180. Is Cirsium palustre a Native of New

Hampshire, 217. Isoëtes, 2; echinospora, 2.

Iva frutescens, 89, 94.

Ivy Poison, 43.

Ivy-poisoning, 105, and its Treatment, 43.

Jasione montana, 15, 20, 199, in Massa-

chusetts, 199. Jewell, H. J., Notes on some Ferns of Franklin County, Maine, 247.

Jones, L. R., Field Meeting of the Vermont Botanical Club, 189; Pogonia affinis in Vermont, 216; Pringle and Frost Herbaria at the University

of Vermont, 171. Josselyn Botanical Society of Maine, Announcement, 110; Annual Meeting,

188; Mid winter Meeting, 62.

Juneus, 60, 170; balticus, 33; brachycarpus, 27, 60; canadensis, var. subcaudatus, 99; filiformis, 2; Greenei, 85; nodosus, 60, var. megacephalus, 170; pelocarpus, 129; scirpoides, 60; subtilis, 129; tenuis, var. anthelatus, 27, var. Williamsii, 99; Torreyi, 170, Leavitt, R. G., Notes on Lycopodium,

and Ellisia Nyctelea in Massachusetts, 170.

Jungermannia barbata, 30, var. attenuata, 210, var. Floerkii, 210, var. lycopodioides, 210, var. quinquedentata, 210; bicrenata, 209; capitata, 209; excisa, 209, var. crispa, 209; Floerkii, 210; incisa, 242; intermedia, 209; lycopodioides, 210; Lyoni, 210; Mildeana, 212; Naumanni, 210; Novae-Caesareae, 212; quinquedentata, 210, 211; Schraderi, 242; ustulata, 208; ventricosa, 242.

Juniper, 34.

Juniperus communis, var. canadensis, 34, 133; Sabina, var. procumbens, 34, 37; virginiana, 133.

Kalmia glauca, 106, 189; latifolia, 131,

Kantia Sullivantii, 207, 212; Trichomanis, 211.

Kennedy, G. G., Maine Coast at Cutler, 23; Plants new to Eastern Massachusetts, 60.

King-Devil Weed, 151, in the Penobscot Valley, 61.

Knight, O. W., "King-devil Weed" in the Penobscot Valley, 61.

Knotgrass, Bird's, 65.

Knowlton, C. H., Occurrence of Empetrum in Franklin County, Maine, 196.

Labrador Tea, 131.

Lady's Slipper, 94, 95, 106.

Lactuca hirsuta, 37; Morssii, 138; scariola, var. integrata, 92.

Lake St. John, Quebec, Two more Rare Plants from, 128.

Laminaria, 175, 177; digitata, var. typica, 177; longicruris, 177; platymeris, 176; stenophylla, 175.

Lamium, 218; album, 218, in Maine, 218. Lapsana communis, 92.

Larch, 35. Large Cranberry, 234.

Larix americana, 106, 189.

Lastraea, 10, 53, 161, 162; dilatata, var. tanacetifolia, 53; Filix mas, 159; patens, 162; tetragona, 162.

Laternea, 135.

Laurel, Mountain, 131, 140, 142, 151.

Lawrence, Massachusetts, Clathrus columnatus in, 134; Volvaria volvacea

in, 3. Leak, 81.

folia, 183; Notice of work, 85.

Lechea major, 28, in New Hampshire, 28; minor, 130.

Ledum groenlandicum, 131; latifolium,

Lee, L. A., Aquilegia canadensis, var. flaviflora in Maine, 169.

Leersia virginica, 93.

Leguminosae, 13.

Leontodon, 249; autumnalis, 39, 40, 189, 249, var. pratensis, 189; hastilis, 250; in America, Early Records of, 39; Newly Introduced Species of Crepis and, 249; palustris, 157. Leonurus Cardiaca, 91.

Lepidium sativum, 32. Lepidozia reptans, 30.

Lepiota americana, 199; cepaestipes, 199; naucina, 199.

Leptobryum pyriforme, 30, 182. Leptopoda brachypoda, 198.

Leskea nervosa, 182; polycarpa, 239.

Leucobryum vulgare, 30.

Leucodon sciuroides, 30, 31, 183. Leucophysalis grandiflora, 190.

Lichens, 172.

Ligustrum vulgare, 17, 74.

Lilium canadense, 89; philadelphicum,

Lily, Cow, 93.

Limnanthemum lacunosum, 76.

Limnorchis, 79; dilatata, 80; fragrans, 80, 81; huronensis, 79, 81; hyperborea, 80, 81; major, 79, 81; media, 79, 80,

Limosella aquatica, var. tenuifolia, 33.

Lindera Benzoin, 132.

Lingonris, 232.

Liochlaena lanceolata, 30.

Liparis liliifolia, 18, 20, 21, 74; Loeselii, 19, 62, in Massachusetts, 62.

List of Bryophytes from the Mt. Greylock Region, 29.

Listera auriculata, 19; convallarioides, 19; cordata, 19, 21.

Lists of New England Plants, Preliminary,- VIII, 15; IX, Polygonaceae, 203; X, Carex, 218.

Litorella lacustris, 129.

Liverwort, 81.

Lobelia cardinalis, 15; Dortmanna, 15, 76; inflata, 15; Kalmii, 15; spicata, 15, 75; syphilitica, 15, 20.

Lobeliaceae, 15.

Local Flora, A New, 202.

Lonicera, 40; caerulea, 24; glauca, 131, sempervirens established in Westford, Massachusetts, 40.

57; Seed Dispersal of Viola rotundi- Lophanthus, 88; scrophulariaefolius, 89, 94.

DECEMBER

Lophocolea heterophylla, 30.

Lophozia barbata, 210, 211; bicrenata, 209; Floerkii, 210, 211; gracilis, 210; lycopodioides, 210, 211, var. Floerkii, 210; Lyoni, 210, 211; Marchica, 211, 212; quinquedentata, 210, 211.

Lousewort, 88. Lupinus perennis, 76. Lychnis chalcedonica, 85.

Lycopodium, 55; chamaecyparissus, 59; clavatum, 57, 101, 102, and its Variety monostachyon, Notes on, 100, var. monostachyon, 57, 58, 100, 101, on Mt. Monadnock, 57; complanatum, 57, 59, 154, extension of Range for the Typical, 154, var. chamaecyparissus, 59, 98, var. flabelliforme, 59; lucidulum, 58, Gemmae of, 58; Notes on, 57; obscurum, 57; Selago, 58, 150; sitchense, 98, on Mt. Washington, Discovery of Comandra livida and, 97; tristachyum, 98.

Lycopus virginicus, 91.

Lygodium, 83; palmatum, 36, 51, 64, 83. Lyngbya aestuarii, 123.

Lysimachia Nummularia, 16; producta, 16, 92, 137; punctata, 16, 62; quadrifolia, 16; stricta, 16; thyrsiflora, 16; vulgaris, 16, 107

Lythrum Hyssopifolia, 89.

Magnolia glauca, 140.

Maiden-hair, False, 54. Maine, Algologist's Vacation in Eastern, 174; Aquilegia canadensis, var. flaviflora in, 169; Aulacomnium heterostichum in, 169; Coast at Cutler, 23; Further Notes on Solanum rostratum and Hieracium praealtum in, 151; Josselyn Botanical Society of, Announcement, 110, Annual Meeting, 188, Midwinter Meeting, 62; Lamium album in, 218; New Stations for Peltandra virginica and Conopholis americana in, 168; Notes on some Ferns of Franklin County, 247; Occurrence of Eleocharis diandra at Brunswick, 1; Occurrence of Empetrum in Franklin County, 196; Seneca Snakeroot in, 133. Mann, R. L. An American Occurrence

of the European Centaurea diffusa,

Maple, River, 130; Silver, 130. Marchantia polymorpha, 30. Marginal Shield-fern, 51.

Margined Sumach, 130.

Marine Cladophoras of New England,

Marsupella, 208; adusta. 208; emarginata, 208, 209, 242; sphacelata, 208;

Sprucei, 208; ustulata, 208.

Massachusetts, Clathrus columnatus in Lawrence, 134; Crepis virens in Eastern, 197; Cuphea procumbens at Andover, 247; Cypripedium arietinum on Mt. Toby, 62; Hemicarpha in Eastern, 82; Hieracium praealtum at Andover, 197; Jasione montana in, 199; Juncus Torreyi and Ellisia Nyctelea in, 170; Liparis Loeselii in, 62; Lonicera sempervirens established in Westford, 40; Plants new to Eastern, 60; Ranunculus abortivus, var. eucyclus in Hyde Park, 152; Rare Plants in Centreville, 245; Some Additions to the Flora of, 166; Some Plants recently found in and around North Easton, 13; Two Noteworthy Carices at Sudbury, 167; Utricularia minor in Holbrook, 42, in Vinnicunnet, 102; Volvaria volvacea in Lawrence, 3.

Mathews, F. S., Notice of work, 201. Matricaria discoidea, 33, 107, 189; ino-

dora, 189.

Meeting of the Josselyn Botanical Society, Annual, 188, Midwinter, 62; Vermont Botanical Club, Field, 189.

Menispermum canadensis, 129.

Mentha aquatica, 106; canadensis, 34; Cardiaca, 100; piperita, 192; rubra, 34. Menyanthes trifoliata, 106.

Merrill, E. D., Notes on North American Grasses, 142; Notes on Sporobolus, 45.

Mertensia maritima, 189. Mesogloia divaricata, 178.

Metcalf, Mrs. R. E., Climbing Fern in

New Hampshire, 83.

Metzgeria conjugata, 30.

Microstylis, 75.; monophyllos, 19; ophioglossoides, 19, 75.

Midwinter Meeting of the Josselyn Botanical Society of Maine, 62.

Milium, 142, 143; Muhlenbergianum, 142, 143; pungens, 142, 143.

Mint, 34. Miscellaneous Notes on New England Ferns,—III, 7, IV, 49, V, 157.

Mistletoe, 35; at Bradford, Vermont,

The Dwarf, 28; Dwarf, 35. Mnium affine, 30; cuspidatum, 30; orthorrhynchum, 182; punctatum, 30, var. elatum, 30; serratum, 30; stellare, 30.

Monadnock, Lycopodium clavatum, var. monostachyon on Mt., 57.

Monarga fistulosa, 91; mollis, 89, 91.

Monostroma latissimum, 175; quaternarium, 175.

Montia fontana, 150.

Moonseed, 129.

Morrell, J. M. H., Lamium album in Maine, 218.

Mosses, 29; Notes on Vermont, 180. Moulton, D. H., Annual Meeting of the Josselyn Society, 188; Josselyn Botanical Society of Maine, Announcement, 110; Midwinter Meeting, 62. Mount Greylock, Pogonatum capillare

on, 83.

Mount Greylock Region, Bryophytes of the,—II. 238; List of Bryophytes from the, 29.

Mount Monadnock, Lycopodium clavatum, var. monostachyon on, 57.

Mount Toby, Massachusetts, Cypripe-

dium arietinum on, 62.

Mount Washington, Discovery of Comandra livida and Lycopodium sitchense

Mountain Cranberry, 231, 232; Laurel. 131, 140, 142, 151; Pink, 131; Spleenwort, 54.

Mouse-ear Hawkweed, 151.

Muhlenbergia aspericaulis, 46; capillaris, 143; dumosa, 46; expansa, 143; trichopodes, 143.

Musci, 30, 238. Mushroom for the market, A New, 199. Mushrooms, Certain eaters of, 77.

Mutinus caninus, 134. Myosotis arvensis, 91.

Myriophyllum, 105; ambiguum, 105, var. limosum, 105.

Narcissus poeticus, 84.

Nardia adusta, 208; haematosticta, 209; hyalina, 209.

Native Flora, Destruction of our, 200; Protection of our, 139.

Neckera pennata, 30. Necklace Poplar, 133.

Nephrodium, 9, 10, 158-166; acrostichoides, 159, 162; Boottii, 10, 52; Clintonianum, 10-12; cristatum, 10, 52, 53, 165, var. Clintonianum, 52, var. Slossonae, 52, × marginale, 10, 11, 51; dilatatum, 53; Filix-mas, 158, 159, 162, 163, 165; fragrans, 10; Goldieanum, 10, 52; marginale, 9-12, 51, 165, 245, Cristate form of, 244, forma Davenportii, 245; molle, 161; noveboracense, 10, 165; simulatum, 10, 84; Some Further observations on, 158; spinulosum, 9, 10, 12, 53, var. dilatatum, 53, var. intermedium, 53; The-

lypteris, 10; unitum, 161. New Bedford, Rare Plants about, 106. New England Aspidieae, 7; Chilian Empetrum in, 147; Evergreen Ferns of, 49; Ferns, Miscellaneous Notes on, III, 7, IV, 49, V, 157; Handbook of the Trees of, 40; Hepaticae, Notes on, 207; Ma-rine Cladophoras of, 111; Plants, Preliminary lists of, VIII, 15, IX, Polygonaceae, 203, X, Carex, 218; Polygonums of the Section Avicularia, 65; Rediscovery of Phaseolus perennis in, 13; Valerianella in, 195; Veronica Chamaedrys in, 107.

New Hampshire, Calluna vulgaris in Southern, 200; Climbing Fern in, 83; Is Cirsium palustre a Native of, 217; Lechea major in, 28; Plants in the Middle Connecticut Valley, Some Vermont and, I, 129; Some Plants

of Intervale, 61.

New Hypericums of the Adpressum group, Two, 135; Jersey Tea, 130; Local Flora, 202; Mushroom for the Market, 199; Popular Field Book, 201; Station for Polypodium vulgare, var. cambricum, 197; Stations for Peltandra virginica and Conopholis americana in Maine, 168.

North America, Cotton-grass new to, 82. North American Grasses, Notes on, 142. North Easton, Massachusetts, Some Plants found in and around, 13.

Northeastern Veronicas, Two, 191. Norton, A. H., New Stations for Peltandra virginica and Conopholis americana in Maine, 168.

Norway Pine, 133.

Notable Plant Stations in Rhode Island, Some, 108.

Note upon Recent Treatment of Habenaria hyperborea and its Allies, 79.

Notes on Lycopodium, 57; Lycopodium clavatum and its Variety monostachyon, 100; New England Ferns, Miscellaneous, III, IV, V, 7, 49, 157; New England Hepaticae, 207; North American Grasses, 142; Solanum rostra- Pastinaca sativa, 188. tum and Hieracium praealtum in, Maine, Further, 151; Some Ferns of Pease, A. S., Hieracium praealtum at Franklin County, Maine, 247; Sporobolus, 45; the Blue-berried Huckleberry, 14; the Flora of Connecticut, 36; Vermont Mosses, 18o.

Noteworthy Carices at Sudbury, Massachusetts, Two, 167; Plants of Connecticut, 84. 98; Plants of Southeastern Connecticut,- III, 26.

Nye, H. A., Blooming of Hepaticas, 127. Nymphaea odorata, var. minor, 76, 103;

reniformis, 27; tuberosa, 27. Nyssa sylvatica, 91, 131.

Oak, 93; Black, 132; Island and its Flora, 87; Scrub Black, 133; White, 132; Yellow, 132.

Observations on Nephrodium, Some

Further, 158.

Occurrence of Eleocharis diandra at Brunswick, Maine, 1; of Empetrum in Franklin County, Maine, 196; of the European Centaurea diffusa, American, 249.

Odontoschisma denudatum, 211.

Oleaceae, 17.

Onion, 81.

Onoclea sensibilis, 157, forma obtusilobata, 36; Struthiopteris, 157.

Orange Hawkweed, 62, 151.

Orchidaceae, 18.

Orchis rotundifolia, 19, 21, 22; spectabilis, 19, 22, 93.

Orthotrichum sordidum, 181; strangulatum, 30.

Oryzopsis canadensis, 142, 143; juncea, 142, 143; parviflora, 142, 143.

Newly Introduced Species of Crepis and Osmorrhiza, 88, 153, 154; brevistylis, Leontodon, 249.

153; longistylis, 81, 153; New to Eastern America, 153; obtusa, 154.

Osmunda, 83; regalis, 83. Our Chokeberries, 55. Outlines of Botany, 85. Oxyria digyna, 203.

Paint-brush, Devil's, 62.

Panicularia borealis, 146; Davyi, 145;

fluitans, 146; Torreyana, 146.

Panicum boreale, 89; dichotomum, 38; firmum, 142, 143; lanuginosum, 38, 90; longifolium, 38; macrocarpon, 38, 89; microcarpon, 39; sphaerocarpon, 39; stipitatum, 26; unciphyllum, 38,

Paronychia argyrocoma, 61.

Parsnip, 188; Virulence of the Wild, 188. Paspalum Muhlenbergii, 26.

Pea, Hoary, 249.

Andover, Massachusetts, 197; Two new Stations for Arceuthobium, 249. Pedicularis canadensis, 88, 93.

Pellaea atropurpurea, 54.

Peltandra, 169; virginica, 106, 168, and Pleocnemia, 162. Conopholis americana in Maine, New Stations for, 168.

Penobscot Valley, "King-devil Weed"

in, 61.

Peppergrass, 32. Pepperidge, 131.

Peridermium abietinum, 35.

Persistence of the Button-bush, Remark-

Petasites palmata, 128.

Pfaff, F., Ivy Poisoning and its Treatment, 43.

Phaseolus, 13; perennis, 13, 99, in New England, Rediscovery of, 13

Philonotis fontana, 30. Phleum alpinum, 150.

Phragmites, 93; communis, 93.

Phryma Leptostachya, 90. Physcomitrium turbinatum, 30. Physostegia virginiana, 190.

Picea alba, 24, 189; rubra, 28.

Pignut, 132.

Pine, 74; Norway, 133; White, 93. Pink, Mountain, 131; Swamp, 131.

Pink-spored Agarics, 6.

Pinus resinosa, 133. Plagiothecium denticulatum, 30, 241; Muellerianum, 182; silvaticum, 241; turfaceum, 30, 182, 241.

Plant Stations in Rhode Island, Some

Notable, 198.

Plantago borealis, 24, 25; decipiens, 25,

89; maritima, 25.
Plants about New Bedford, Rare, 106; from Lake St. John, Quebec, Two more Rare, 128; from Prince Edward Island, Some, 31; in Centreville. Massachusetts, Rare, 245; in the Middle Connecticut Valley, Some Vermont and New Hampshire, - I, 129; mentioned in Young's Chronicles, Concerning the, 81; New to Eastern Massachusetts, 60; of Connecticut, noteworthy, 84, 98; of Intervale, New Hampshire, Some, 61; of Southeastern Connecticut, Noteworthy, — III, 26; of the Blue Hills Reservation, A few, 74; Preliminary Lists of New England, VIII, 15, IX, Polygonaceae, 203, X, Carex, 218; recently found in and around North Easton, Massachusetts, Some, 13.

Platanthera convallariaefolia, 79; dilatata, 79; hyperborea, 79; leucostachys,

Platanus occidentalis, 132.

Plectonema Battersii, 177.

Plum, 81.

Plumaria elegans, 176.

Plumbaginaceae, 16.

Poa, 25; alpina, 25; ampla, 145; arachnifera, var. glabrata, 144; autumnalis, 145; brachyphylla, 145; brevifolia, 145; cuspidata, 145; debilis, 39; elongata, 146; glumaris, 33; Leckenbyi, 145; nevadensis, 145; nitida, 144; pratensis, 25, var. domestica, 25; pungens, 145; Torreyana, 146.

Pogonatum alpinum, 30, 31; capillare, 83, 240, 241, on Mt. Greylock, 83;

urnigerum, 30, 31.

Pogonia affinis, 19, 22, 216, in Vermont, 216; ophioglossoides, 19, 42; pendula, 2, 19, 21, 22, 61, Elusive character of, 2; verticillata, 19, 74, 217.

Poison Ivy, 43; Sumach, 43, 130. Poisoning and its Treatment, Ivy, 43; by Cypripedium spectabile in Vermont,

Some Cases of, 94.

Polygala Nuttallii, 85; paucifolia, 74, 76; polygama, 74, 75, 249; Senega, 133. Polygonaceae, 203.

Polygonella articulata, 61, 204.

Polygonum, 65; acre, 204, var. lepto-stachyum, 204; amphibium, 104, 105, 204, Biological Relations of Polygonum Hartwrightii to, 104, var. Hartwrightii, 105, 204; arenarium, 66, 70, 205; arifolium, 204; aviculare, 65, 66, 69, 72, 73, 204, 205, var. angustissimum, 70, 73, var. erectum, 71, var. laxum, 70, var. littorale, 69, 73, 91, 204, 205, var. neglectum, 70, var. vegetum, 70, 73; Bellardi, 69, 72, 73. 205; camporum; 72; Careyi, 204; cilinode, 204; Convolvulus, 204; cuspidatum, 204, 205; Douglasii, 73, 204, 205; dumetorum, 204; erectum, 71, 72, 73, 204; exsertum, 68, 73, 204; Fowleri, 67, 68, 73, 204; glaucum, 67; Hartwrightii, 104, 105, to Polygonum amphibium, Biological Relation of, 104; Hydropiper, 204; hydropiperoides, 204, var. strigosum, 204; lapathifolium, 204, var. incanum, 204; littorale, 70; marinum, var. roseum, 67; maritimum, 66, 67, 68, 73, 91, 204, var. littorale, 70; Muhlenbergii, 204; neglectum, 70; orientale, 204, 205; pennsylvanicum, 68 204; Persicaria, 204; prolificum, 68, 69, 72, 73, 91, 204; Raii, 33, 67, 88; ramosissimum, 69, 71, 73, 204, forma atlanticum, 72, 204, var. prolificum, 68; Rayi, 67, 68; Roberti, 67, 68, 73;

rurivagum, 69; sagittatum, 204; scandens, 204; tenue, 73, 204; virginianum, 199, 204; viviparum, 204.

Polygonums of the Section Avicularia,

New England, 65.

Polypodium, 165, 247; vulgare, 51, 245, var. auritum, 247, var. cambricum, 51, 197, 247, New station for, 197.

Polypody, Common, 51.

Polysiphonia Olneyi, 178; violacea, 177. Polystichum, 8, 9, 54, 159, 162-164; acrostichoides, 10, 53; aculeatum, var. Braunii, 10, 54; Lonchitis, 8.

Polytrichum commune, 30; juniperinum, 30; ohioense, 30; piliferum, 30.

Pontederia cordata, 168. Poplar, Necklace, 133.

Populus deltoides, 133. Porella platyphylla, 30.

Porotrichum alleghaniense, 183.

Porphyra amplissima, 177; miniata, 177. Potentilla Anserina, 33, 89; canadensis, 91; palustris, 189; tridentata, 38.

Preisselbeere, 232.

Preissia commutata, 30.

Preliminary Lists of New England Plants, -VIII, 15, IX, Polygonaceae, 203, X, Carex, 218.

Prenanthes, 89; alba, 89; trifoliolata, 37, 89, 92.

Prickly Ash, 130.

Primula farinosa, 16, 150; mistassinica,

Primulaceae, 16, 213.

Prince, F. C., Some Plants of Intervale, Remarkable Persistence of the Button-

New Hampshire, 61. bush, 243.
Prince Edward Island, Some Plants Remarks on Volvaria, 5. from, 31.

Pringle and Frost Herbaria at the Uni-

versity of Vermont, 171. Protection against Rhus-poisoning, Sal-

eratus as a, 105; of our Native Flora, 139.

Prunus pumila, 130. Ptilidium ciliare, 30.

Puccinellia angustata, 27.

Purple Cliff-Brake, 54.

Pycnanthemum linifolium, 75.

Pylaisia intricata, 30; velutina, 183.

Pyrus arbutifolia, 55, 56, 131, var. erythrocarpa, 91, var. melanocarpa, 38, 91.

Quebec, Two more Rare Plants from Lake St. John, 128.

Quercus acuminata, 167; alba, 132; ilicifolia, 38, 133; Muhlenbergii, 190; prinoides, 133; velutina, 132.

Racomitrium aciculare, 30, 31; Sudeticum, 241.

Radula complanata, 242; obconica, 207, 213.

Ragwort, Tansy, 34.

Rand, E. L., Hemicarpha in Eastern Massachusetts, 82; Staminate Plant of Antennaria Parlinii, 152.

Range for the Typical Lycopodium complanatum, Extension of, 154.

Ranunculus, 105; abortivus, 88, var. eucyclus, 152, in Hyde Park, Massachusetts, 152; acris, 32, var. Steveni, 32; allegheniensis, 88, 92; aquatilis, var. tricophyllus, 76; Cymbalaria, 33; Flammula, var. reptans, 129; multi-fidus, 105, var. terrestris, 105; recurvatus, 76, 88; repens, 92.

Raphidostegium recurvans, 241.

Rare Plants about New Bedford, 106; from Lake St. John, Quebec, Two more, 128; in Centreville, Massachusetts, 245.

Recent Treatment of Habenaria hyperborea and its American Allies, Note

upon, 79.

Records of Leontodon in America, Early,

Red Cedar, 133; Chokeberry, 55, 131. Rediscovery of Phaseolus perennis in New England, 13.

Reed, Sea-sand, 33.

Relation of Polygonum Hartwrightii to P. amphibum, Biological, 104.

Reservation, A Few Plants of the Blue Hills, 74.

Rhinanthus Crista-galli, 108.

Rhode Island, Some Notable Plant Stations in, 198.

Rhododendron, 141; canescens, 131; nudiflorum, 100.

Rhodophyllis dichotoma, 176.

Rhus, 97; copallina, 130; glabra, 130; Toxicodendron, 43, 95; venenata, 43, 95, 105, 130.

Rhus-poisoning, Saleratus as a Protec-

tion against, 105.

Rhynchostegium rusciforme, 241; serratulum, 183.

Riccia crystallina, 207.

Rich, W. P., Juncus Torreyi and Ellisia Nyctelea in Massachusetts, 170; Oak Island and its Flora, 87; Preliminary Lists of New England Plants,-IX, Polygonaceae, 203.

River Maple, 130; Rose, 130.

Robinia hispida, 198.

Robinson, B. L., Further Stations for Veronica Chamaedrys, 152; Lechea major in New Hampshire, 28; New England Polygonums of the Section Avicularia, 65; Protection of Our Native Flora, 139; Two New Hypericums of the Adpressum Group, 135; Veronica Chamaedrys in New England, 107.

Robinson, J., Concerning the Plants mentioned in Young's Chronicles, 81.

Rock Cranberry, 231.

Rosa blanda, 130; carolina, 131; lucida, 189, var., 34.

Rose, River, 130; Smooth, 130; Swamp,

131.

Round-leaved Dogwood, 131.

Rubus, 64, 97; argutus, 91; Chamaemorus, 97, 189; nigrobaccus, 91; sp.,

Rumex Acetosa, 204; Acetosella, 204; altissimus, 204; Brittanica, 204; conglomeratus, 205; crispus, 204; hastatulus, 204, 205; maritimus, 205; obtusifolius, 204; Patientia, 204; persicarioides, 205; salicifolius, 205; sanguineus, 205; verticillatus, 205.

Sabbatia chloroides, 103, 107; stellaris,

Sagenia, 162.

Sagina nodosa, 24, var. pubescens, 24, 26; procumbens, 150.

Sagittaria, 2, 168; rigida, 39. St. John, Quebec, Two more Rare Plants from Lake, 128.

Saleratus as a Protection against Rhuspoisoning, 105.

Salix longifolia, 99. Salsola Kali, 33.

Samolus Valerandi, var. americanus, 16.

Sand Cherry, 130.

Sandford, S. N. F., Cut-leaved Cherry Birch, 83. Sanicula gregaria, 89, 91; marylandica,

88. Santalaceae, 18.

Sarcoscyphus adustus, 208; Ehrharti; pigmaeus, 208; Sprucei, 208; ustulatus, 208.

Sargent, C. S., Some Additions to the Flora of Massachusetts, 166.

Sarracenia purpurea, 106.

Sassafras officinale, 132.

Scabiosa australis, 106, 107.

Scapania, 212; curta, 212, 213; nemo-

rosa, 242; undulata, 30, 31, var puipurea, 31.

Scheuchzeria palustris, 189.

Schistostega, 182; osmundacea, 182. Scirpus lacustris, 102; pungens, 33.

Scorgie, A. M., Jasione montana in Massachusetts, 199.

Scrophularia marylandica, 89, 94.

Scrub Black Oak, 133.

Scutellaria, 137; Churchilliana, 138; galericulata, 137, 138; lateriflora, 137, 138.

Sea Sand Reed, 33.

Seaside Golden-rod, 89.

Section Avicularia, New England Polygonums of the, 65.

Sedum acre, 199; Rhodiola, 189. Seed Dispersal of Viola rotundifolia, 183. Seed-throwing of Viola, 230.

Selaginella, 55.

Seneca Snakeroot in Maine, 133.

Senecio Jacobaea, 34; Robbinsii, 189; sylvaticus, 34.

Seventh Annual Winter Meeting of the Vermont Botanical Club, 64.

Shagbark, 132.

Shaw, E. L., Carex aurea in Connecticut, 168; New Station for Polypodium vulgare, var. cambricum, 197

Sheldon, J. L., Notes on the Blue-berried Huckleberry, 14.

Shield-fern, Marginal, 51.

Sibbaldia, 98.

Silene Armeria. 85. Silsbee, F. H., Volvaria volvacea in Lawrence, Massachusetts, 3.

Silver Maple, 130.

Sisyrinchium angustifolium, 91; Bermudiana, 91.

Skullcap, An Anomalous, 137.

Small Cranberry, 235. Smilacina stellata, 34, 76.

Smith, A. M., Pogonatum capillare on Mt. Greylock, 83.

Smooth Rose, 130; Sumach, 130.

Snakeroot, 88; in Maine, Seneca, 133. Snowflake, 218.

Society, Annual Meeting of the Josselyn Botanical, 62, 110, 188; for the Protection of Native Plants, 139.

Solanum, 13; carolinense, 13; Dulcamara, 91; Lycopersicum, 28; rostratum, 14, 151, and Hieracium praealtum in

Maine, Further Notes on, 151. Solidago, 89; altissima, 92; arguta, 89;

asperula, 89, 92; canadensis, var. procera, 92; Elliottii, 38; sempervirens, 89, 92; sp., 92; uliginosa, 38.

Some Additions to the Flora of Massachusetts, 166; Cases of Poisoning by Cypripedium spectable in Vermont, 94; further Observations on Nephrodium, 158; Notable Plant Stations in Rhode Island, 198; Noteworthy Plants of Connecticut, 98; Plants from Prince Elward Island, 31; Plants of Intervale, New Hampshire, 61; Plants recently found in and around North Easton, Massachusetts, 13; Vermont and New Hampshire Plants in the Middle Connecticut Valley,- I, 129.

Sonchus asper, 92.

Sorrel, 81.

Southeastern Connecticut, Noteworthy Plants of,- III, 26.

Southern New Hampshire, Calluna vulgaris in, 200.

Spartina caespitosa, 27; juncea, 27 Species of Crepis and Leontodon Newly

Introduced, 249. Specularia perfoliata, 15.

Speedwell, 107.

Spergularia borealis, 33.

Sphagnum, 42, 74, 75, 212, 238; acutifolium, 30, 242; cuspidatum, 241; cymbifolium, 30; intermedium, 242; medium, 242; recurvum, 242; rigidum, 242; squarrosum, 30; tenerum, 242.

Spice-bush, 132.

Spiraea japonica, 28; prunifolia, 28. Spiranthes cernua, 19, 76; gracilis, 19; latifolia, 19; praecox, 19, 20; Romanzoffiana, 19, 22; simplex, 19. Spleenwort, Ebony, 54; Green, 54;

Mountain, 54.

Spongomorpha, 113-115, 119; Hystrix, 116, 117; Sonderi, 116, 117; spinescens, 117.

Sporobolus aristatus, 48; aspericaulis, 46; brevifolius, 46, 47, 49; cuspidatus, 47; depauperatus, 45, 46, var. filiformis, 46; ejuncidus, 48, 49; filiformis, 47, 48; gracilis, 48, 49; gracillimus, 47; junceus, 48, 49; Notes on, 45; Richardsonii, 45-47; simplex, 47, 48, var. thermale, 48.

Spruce, 93; Black, 35; White, 35.

Stachys palustris, 85, 103.

Staminate Plant of Antennaria Parlinii.

Staphylea trifolia, 130, 190.

Statice Limonium, var. caroliniana, 16. Station for Polypodium vulgare, var. cambricum, New, 197

Stations for Arceuthobium, Two New, 249; for Peltandra virginica and Cono- Triglochin maritima, 89.

pholis americana in Maine, New, 168; for Veronica Chamaedrys, Further,

Stature attained by Euphrasia Oakesii, 108.

Steironema ciliatum, 16, 88; lanceolatum, 16.

Stinking Willie, 34.

Stipa canadensis, 142, 143; expansa, 143; juncea, 142, 143.

Strawberry, 81, 128.

Suaeda, 33.

Subularia aquatica, 129.

Massachusetts, Two Note-Sudbury, worthy Carices at, 167. Sumach, Dwarf, 130; Margined, 130;

Poison, 43, 130; Smooth, 130.

Summer Grape, 130.

Swamp Pink, 131; Rose, 131. Sweet Coltsfoot, 128; Fern, 40.

Symphoricarpos vulgaris, 106. Symphytum asperrimum, 84.

Syringa vulgaris, 17, 20.

Tanacetum vulgare, 92.

Tansy, 32; Dwarf, 32; Ragwort, 34. Taraxacum Dens-leonis, 155; erythrospermum, 92, 155-157; officinale, 155-157, var. palustre, 155, 157; palustre, 155-157, in America. 155.

Tea, Labrador, 131; New Jersey, 130.

Tectaria, 164.

Tetraphis pellucida, 30.

Tetraspora, 175.

Teucrium canadense, 89.

Thelia asprella, 182.

Thuidium delicatulum, 30; microphyllum, 181; paludosum, 182; scitum,

Thuya occidentalis, 37.

Tillaea, 35; aquatica, 35; simplex, 35, 36; Vaillantii, 35, 36. Tipularia discolor, 19, 21, 22.

Toadstools, 78.

Toxicodendric acid, 43.

Toxicodendrol, 43, 44. Trailing Arbutus, 140.

Treatment of Habenaria hyperborea and its Allies, Note upon Recent, 70.

Trees of New England, Handbook of the, 40.

Trichochloa expansa, 143.

Trichocolea tomentella, 243.

Tricholoma, 77; personatum, 199; portentosum, 77.

Trientalis americana, 16.

Trillium, 128; erythrocarpum, 106; grandiflorum, 198.

Triosteum aurantiacum, 88, 92; perfoliatum, 75, 92, 195.

Trisetum subspicatum, 150. Trumpet Honeysuckle, 40.

Two more Rare Plants from Lake St. John, Quebec, 128; New Hypericums of the Adpressum Group, 135; New Stations for Arceuthobium, 249; Northeastern Veronicas, 191; Noteworthy Carices at Sudbury, Massachusetts, 167.

Typical Lycopodium complanatum, Extension of Range for the, 154.

Tyttebaer, 232.

Ulota crispa, 30; crispula, 30; Hutchinsiae, 30; Ludwigii, 30; phyllantha, 24.

University of Vermont, Pringle and Frost Herbaria at, 171.

Urachne brevicaudata, 142, 143; cana-

densis, 142.

Utricularia, 42, 102, 103; biflora, 27; cornuta, 76, 102; inflata, 76, 102; minor, 2, 42, 102, 103, in Holbrook, Massachusetts, 42, in Vinnicunnet, Massachusetts, 102; purpurea, 76, 102; resupinata, 27; vulgaris, 76, 102.

Uvularia perfoliata, 88; sessilifolia, 88.

Vacation in Eastern Maine, Algologist's

Vaccinium, 14, 232; buxifolium, 233, 234; macrocarpon, 42, 234-237; Oxycoccus, 106, 235-237, var. intermedium, 236, 237; stamineum, 246; vacillans, 131; Vitis-Idaea, 34, 189, 232-234, 237; American Variety of, 231, var., 233, 234, var. major, 233, 237, var. microphyllum, 234, var. minor, 233, 234, 237, var. pumilum, 234.

Valerianella, 195, 196; in New England, 195; radiata, 195, 196, var. leiocarpa,

Variations and Distribution of American Cranberries, 231; of Glaux in America,

Veratrum viride, 74, 76.

Verbena, 246; hastata, 93, 246, forma

rosea, 245.

Vermont, Anagallis arvensis and caerulea in, 185; and New Hampshire Plants in the Middle Connecticut Valley, Some, - I, 129; Bird Club, 138; Botanical Club, 138, Field Meeting of, 189, Seventh Annual Winter Meeting

of, 64; Dwarf Mistletoe at Bradford, 28; Mosses, Notes on, 180; Pogonia affinis in, 216; Pringle and Frost Herbaria at the University of, 171; some Cases of Poisoning by Cypripedum spectabile in, 94.

Veronica, 108, 192, 193; alpina, 194; americana, 192, 193; Anagallis, 192; Beccabunga, 81, 191-193; Chamaedrys, 107, 108, 152, Further Stations for, 152, in New England, 107; humifusa, 194; intermedia, 192; serpyllifolia, 193, 194, var. alpina, 194, var. borealis, 194, var. humifusa, 194, 195.

Veronicas, Two Northeastern, 191. Viburnum, 89, 108; dentatum, 89; Opulus, 131; pubescens, 190.

Vilfa, 49; cuspidata, 47; depauperata, 45, 46, var. filiformis, 47, 48; gracilis, 47-49; gracillima, 47; juncea, 49; Richardsonis, 46; squarrosa, 45, 46; subsetacea, 48, 49.

Vinca minor, 17.

Vincetoxicum nigrum, 17, 84.

Vinnicunnet, Massachusetts, Utricularia

minor in, 102.

Viola, 64; alpestris, 230; canina, 183; ovata, 39; pubescens, 76, 88; rotundifolia, 183, 184, 230, Seed Dispersal of, 183; Seed-throwing of, 230.

Violet, 230; White, 128; Yellow, 88.

Virginia Creeper, 141.

Virulence of the Wild Parsnip, 188.

aestivalis, 92; bicolor, Labrusca, 130.

Vitis-Idaea Vitis-Idaea, 234.

Volvaria, 3, 5-7; bombycina, 6; Loveiana, 7; Remarks on, 5; speciosa, 6, volvacea, 4, 5, 7, in Lawrence, Massachusetts, 3.

Walking Leaf, 55.

Washingtonia, 153; obtusa, 153, 154.

Water Cress, 81, 191.

Webera elongata, 30; nutans, 30.

Webster, H., Certain Eaters of Mush-rooms, 77; Clathrus columnatus in Lawrence, Massachusetts, 134; Form of the Bitter Boletus, 187; New Mushroom for the Market, 199; Remarks

on Volvaria, 5. Webster, J. R., Crepis virens in Eastern

Massachusetts, 197.

Weed, in the Penobscot Valley, "King; Devil," 61; King Devil, 151.

Westford, Massachusetts, Lonicera sempervirens established in, 40.

White Birch, 83; Oak, 132; Pine, 93 Spruce, 35; Violet, 128.

Wickopy, 152.

Wild Parsnip, Virulence of the, 188.
Williams, E. F., Preliminary Lists of
New England Plants,—VIII, 15; Two

Noteworthy Carices at Sudbury, Massa-

chusetts, 167.

Winter Meeting of the Vermont Botanical Club, Seventh Annual, 64.

Witches' Brooms, 249, 250.

Woodsia ilvensis, 247; obtusa, 157.

Woodwardia angustifolia, 36.

Xanthoxylum americanum, 130. Xyris, 76; caroliniana, 76.

Yarrow, 81.

Rhodora

Yellow Erythronium, 128; Oak, 132; Violet, 88.

Young's Chronicles, Concerning the Plants mentioned in, 81.

Zizia aurea, var. obtusifolia, 37.

### BOTANICAL PUBLICATIONS

SYNOPTICAL FLORA OF NORTH AMERICA, by A. Gray and others. Vol. I. Fascicles 1 and 2. A critical treatment of forty-five families of polypetalæ (*Ranunculaceæ to Polygalaceæ*) 1895-1897. \$5.20.—Gray Herbarium of Harvard University, Cambridge, Mass.

FLORA OF MT. DESERT ISLAND, MAINE, by EDWARD L. RAND and JOHN H. REDFIELD. With a Geological Introduction by WILLIAM MORRIS DAVIS. 1894. And a new map of the Island. 1901. Price \$2.00, post free. — Address EDWARD L. RAND, 53 State Street, Boston, Mass.

NEW ENGLAND WILD FLOWERS AND THEIR SEASONS, by WILLIAM WHITMAN BAILEY. Price 75 cents net.—Preston & ROUNDS Co., Providence, R. I.

ARE YOU INTERESTED IN NATURE STUDY? If you are you will want a copy of Mosses with a Hand-lens, by Dr. A. J. Grout. It is the only book of its kind in the English language. It makes the mosses as easy to study as the flowering plants. Eight full page plates and ninety figures in the text. Price \$1.10 postpaid. Send for sample pages to O. T. Louis, 59 Fifth Avenue, N. Y. City.

WANTED. A copy of the third edition of Gray's Manual of Botany of the Northern United States. Please state condition and price. Address Wm. P. Rich, 300 Massachusetts Avenue, Boston, Mass.

# RARE VERMONT PLANTS. MT. WASHINGTON PLANTS.

SOME THIRTY SPECIES OF CRATAEGUS (type material)
ANTENNARIA, SCIRPUS, RUBUS, ETC.

Quality and quantity guaranteed. Price \$10.00 per hundred, post or express prepaid.

W. W. EGGLESTON,
23 N. Main Street,
Rutland, Vt.

FOR SALE. Plowright's British Uredineae and Ustilagineae, Ellis and Everhart's North American Pyrenomycetes, several numbers of vols. 5-7 of the Journal of Mycology, Plantae Wrigtianae, part I by Asa Gray; Grasses of the Pacific Slope, Vasey; Grasses of the Southwest, Vasey. The last two are Bulletins 12 and 13 of the Department of Agriculture. Also a nearly complete set of the Fern Bulletin.

A. J. GROUT, 360 Lenox R'D, Brooklyn, N. Y.

Advertisements of Nurserymen and Dealers in Botanical and other Scientific Publications are inserted in these pages at the following rates per space of 4 in. w 2-2 in. 1 year \$4.00, 6 months \$2.50.

## THE STANDARD BOTANIES.

- GRAY'S MANUAL OF BOTANY. Sixth edition. Revised and extended westward to the one-hundredth meridian, by Sereno Watson, Curator of the Gray Herbarium, Harvard University, and John M. Coulter, Professor of Botany in Wabash College, assisted by specialists in certain groups. Cloth, 8vo, 828 pages, and 25 plates. Price, \$1.62, postpaid.
- GRAY'S FIELD, FOREST, AND GARDEN BOTANY. New edition. Revised and extended by Professor L. H. Bailey, of Cornell University. Cloth, 8vo, 519 pages. Price, \$1.44, postpaid.
- GRAY'S LESSONS AND MANUAL. Comprising the Lessons in Botany, Revised, and the Revised Manual. Cloth, 8vo, 1,056 pages, and 25 plates. Price, \$2.16, postpaid.
- GRAY'S SCHOOL AND FIELD BOOK OF BOTANY. Comprising the Revised Lessons in Botany, and the Revised Field, Forest, and Garden Botany. The text-book in most common use in the high schools of the United States. Cloth, 8vo, 745 pages. Price, \$1.80, postpaid.

AMERICAN BOOK COMPANY, 93 Summer St., Boston, Mass.

New York.

Cincinnati.

Chicago.

#### HARDY NATIVE AMERICAN PLANTS

and Carolina Mountain Flowers. Many new rare and local varieties.

HARLAN P. KELSEY, Landscape Architect,

Prop. Highlands Nursery, Kawana, North Carolina.

Catalogues.

Tremont Building, Boston.

Consultation and plans for improvement of estates, private and public grounds, suburban properties and resorts.

## THE PLANT WORLD,

An Illustrated Monthly Journal of Popular Botany, Established 1897.

Edited by F. H. Knowlton, Charles Louis Pollard and Cornelius L. Shear.

#### NOW INCLUDES THE ASA GRAY BULLETIN,

Each issue consisting of 20 pages, with a supplement of 8 pages, describing the families of flowering plants in systematic sequence. A copy of the new "Flora of Vermont" free to every new subscriber desiring it. Back volumes of The Asa Gray Bulletin can be furnished in limited quantity. Subscription Price of The Plant World \$1.00 per year. Sample copy free on receipt of 1-cent stamp if Rhodora is mentioned. Address all communications to

The Plant World Co.,

Box 334, Washington, D. C.